



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The QT interval increases with decreasing plasma potassium in the Danish general population

Kanters, Jørgen K.; Graff, Claus; Henriksen, Lise; Petri, Anne Sofie; Christiansen, Michael; Nielsen, Jimmi; Kvetny, Jan; Ellervik, Christina

Publication date:
2015

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Kanters, J. K., Graff, C., Henriksen, L., Petri, A. S., Christiansen, M., Nielsen, J., Kvetny, J., & Ellervik, C. (2015). *The QT interval increases with decreasing plasma potassium in the Danish general population*. Poster presented at Annual Conference of The International Society for Computerized Electrocardiology, ISCE , San Jose, CA, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

The QT interval increases with decreasing plasma potassium in the Danish general population

Jørgen K. Kanters^{1,7}, Claus Graff², Lise Henriksen¹, Anne Sofie Petri¹, Michael Christiansen³, Jimmi Nielsen⁴, Jan Kvetny⁵, Christina Ellervik⁶

¹Laboratory of Experimental Cardiology, University of Copenhagen, Denmark, ²Department of Health Science and Technology, Aalborg University, Denmark, ³Statens Seruminstitut, Copenhagen, Denmark, ⁴Department of Clinical Medicine, Aalborg Psychiatric Hospital, Aalborg, Denmark, ⁵Dept. of Clinical Biochemistry, Naestved Sygehus, Denmark, ⁶Department of Laboratory Medicine, Boston Children's Hospital, Boston, MA, USA, ⁷Department of Cardiology, Herlev & Gentofte University Hospital, Copenhagen, Denmark.



It is a well-known fact that during hypokalemia, the corrected QT interval prolongs and the T-waves flattens and may even get humps, During hyperkalemia the corrected QT interval shortens and T-waves get higher amplitudes and become more peaked. However the magnitude of these repolarization changes during physiological variations of Se-K is unknown.

Methods

Subjects were enrolled from the Danish General Suburban Population Study (GESUS). GESUS is a cross-sectional study of the adult suburban general population in Naestved municipality (70 km south of Copenhagen) initiated in January 2010.



The Full Cohort of participants having an electrocardiogram (ECG) recorded consisted of 9914 subjects. The (assumed) Healthy group excluded participants with self-reported history of myocardial infarction/angina pectoris, history of cerebrovascular incident, diabetes mellitus, Hba1c>42, estimated glomerular filtration rate lower than gender- and age-adjusted 5% percentile, intake of any cardiac medication including diuretics, diabetes or insulin injections, consisted of 6314 patients.

ECGs were recorded on a digital ECG recorder (MAC5000, GE-Medical Systems, Milwaukee, WI, USA), at a sampling rate of 500 Hz. Each 10-second ECG was used to form a median beat in the recorded leads using MUSE/Interval Editor software (GE Health care, Milwaukee, WI, USA), where representative beat waveforms were constructed for each lead after baseline removal and then superimposed to construct a superlead. QT measurements were made automatically using the 12SL-algorithm (12SL, GE-Medical Systems, Milwaukee, WI, USA). QT intervals were corrected for heart rate with Fridericia's formula ($QTcF = (QT/RR)^{1/3}$).

Results

Mean±SD		Full Cohort	Assumed Non-Healthy	Assumed Healthy
<i>Demographics</i>				
n		8779	3438	5341
age	y	56.45±13.53	64.11±11.62*	51.52±12.32
male	%	45.8%	47.1% (p=0.052)	45.0%
<i>ECG parameters</i>				
QTcF	ms	416.5±20.4	420.5±21.8*	413.9±19.0
QTcB	ms	420.9±24.5	427.7±25.2*	416.4±22.9
RR	ms	956.0±163.9	918.6±159.8*	980.2±162.0
<i>Clinical Chemistry</i>				
Se-Potassium	mM	3.883±0.297	3.894±0.342*	3.876±0.264
Se-Sodium	mM	140.70±2.24	140.52±2.52*	140.82±2.03
Creatinine	μM	77.39±19.42	80.94±25.45*	75.14±13.83

Demographics of the Assumed healthy GESUS population. *p<0.05 vs. Healthy



QTcF (ms)	Univariate α	95% CI	Multivariate α	95% CI
age (y)	0.21*	[0.17; 0.25]	0.26*	[0.21; 0.30]
gender (M vs. F)	-7.4*	[-8.4; -6.4]	-6.7*	[-7.7; -5.7]
Se-Potassium (mM)	-8.4*	[-10.3; -6.4]	-8.3*	[-10.2; -6.3]
Se-Sodium (mM)	0.12	[-0.12; 0.38]		
Creatinine (μM)	-0.17	[-0.21; -0.14]		
Intercept			436*	[429; 443]
<i>T-wave Amplitude V5 (μV)</i>				
age (y)	-1.8*	[-2.2; -1.4]	-2.3*	[-2.7; -1.9]
gender (M vs. F)	100*	[90;110]	83*	[95; 120]
Se-Potassium (mM)	103*	[85;123]	83*	[77;115]
Se-Sodium (mM)	-0.09	[-2.6; 1.2]		
Creatinine (μM)	2.0*	[1.6; 2.4]	-0.9*	[-1.4;-0.5]
Intercept			161*	[85; 236]

Univariate and Multivariate linear regression of QTcF and T-wave amplitude in lead V5 (Assumed healthy GESUS population). *p<0.05

Conclusion

- T-wave amplitude decreases **103 μV per mM** decrease in potassium
- QTcF interval increases with **8.4 ms per mM** decrease in potassium
- Since FDA has a threshold of concern of $\Delta QTc=5$ ms, it seems necessary to take plasma levels of potassium into account when analyzing data in studies with drug-induced repolarization changes